

For example, take two hydrogen atoms,  $A$  and  $B$ , and consider the states  $\psi_A \pm \psi_B$ .

## TIGHT BINDING MODEL


### Lecture 20

*A.H. Harker*  
Physics and Astronomy  
UCL


#### 7.6 The tight-binding model

##### 7.6.1 Overview


For materials which are formed from closed-shell atoms or ions, or even covalent solids, the free electron model seems inappropriate. In the tight-binding model, we imagine how the wavefunctions of atoms or ions will interact as we bring them together.



The symmetric (+) form has more screening charge between the nuclei, and has lower energy.



When more atoms are brought together, the degeneracies are further split - to form bands ranging from fully bonding to fully antibonding. Different orbitals can lead to band overlap.



## 7.6.2 Tight-binding theory

Consider an element with one atom per unit cell, and suppose that each atom has only one valence orbital,  $\phi(\mathbf{r})$ . Then we can make a wavefunction of Bloch form by forming

$$\psi_{\mathbf{k}}(\mathbf{r}) = N^{-1/2} \sum_m \exp(i\mathbf{k} \cdot \mathbf{R}_m) \phi(\mathbf{r} - \mathbf{R}_m).$$

Confirm that this is a Bloch function. If  $\mathbf{T}$  is a translation vector:

$$\begin{aligned} \psi_{\mathbf{k}}(\mathbf{r} + \mathbf{T}) &= N^{-1/2} \sum_m \exp(i\mathbf{k} \cdot \mathbf{R}_m) \phi(\mathbf{r} - \mathbf{R}_m + \mathbf{T}) \\ &= N^{-1/2} \exp(i\mathbf{k} \cdot \mathbf{T}) \sum_m \exp(i\mathbf{k} \cdot (\mathbf{R}_m - \mathbf{T})) \phi(\mathbf{r} - (\mathbf{R}_m - \mathbf{T})) \\ &= \exp(i\mathbf{k} \cdot \mathbf{T}) \psi_{\mathbf{k}}(\mathbf{r}) \end{aligned}$$

because if  $\mathbf{R}_m$  is a lattice vector, so is  $\mathbf{R}_m - \mathbf{T}$ .

Find the expectation energy of the Hamiltonian:

$$\langle \mathbf{k} | \mathcal{H} | \mathbf{k} \rangle = N^{-1} \sum_m \sum_n \exp(i\mathbf{k} \cdot (\mathbf{R}_n - \mathbf{R}_m)) \langle \phi_m | \mathcal{H} | \phi_n \rangle$$

where  $\phi_m = \phi(\mathbf{r} - \mathbf{R}_m)$ . Now  $\langle \phi_m | \mathcal{H} | \phi_n \rangle$  will be large if  $n$  and  $m$  are the same atomic site, or nearest neighbours, but will decrease rapidly with separation. Write

$$\begin{aligned} \langle \phi_n | \mathcal{H} | \phi_n \rangle &= -\alpha, \\ \langle \phi_m | \mathcal{H} | \phi_n \rangle &= -\gamma \text{ if } n \text{ and } m \text{ are nearest neighbours,} \\ \langle \phi_m | \mathcal{H} | \phi_n \rangle &= 0 \text{ otherwise.} \end{aligned}$$

Then

$$E_{\mathbf{k}} = \langle \mathbf{k} | \mathcal{H} | \mathbf{k} \rangle = -\alpha - \gamma \sum_n \exp(i\mathbf{k} \cdot \mathbf{R}_n),$$

where the sum is over nearest neighbours only, and  $\mathbf{R}_n$  is a vector joining an atom to its nearest neighbours. For example, in two-dimensional square lattice we have

$$\{\mathbf{R}_n\} = \{(a, 0), (-a, 0), (0, a), (0, -a)\}$$

so that if  $\mathbf{k} = (k_x, k_y)$

$$E_{\mathbf{k}} = -\alpha - 2\gamma(\cos(k_x a) + \cos(k_y a)).$$

Clearly, as  $\cos$  ranges between  $-1$  and  $1$   $E_{\mathbf{k}}$  ranges between  $-\alpha - 4\gamma$  and  $-\alpha + 4\gamma$ , giving a band width of  $8\gamma$ . Near  $\mathbf{k} = 0$  we can expand the  $\cos$  functions as

$$\cos \theta \approx 1 - \frac{1}{2}\theta^2,$$

so

$$\begin{aligned} E_{\mathbf{k}} &\approx -\alpha - 2\gamma\left(1 - \frac{1}{2}k_x^2 a^2 + 1 - \frac{1}{2}k_y^2 a^2\right) \\ &= -\alpha - 4\gamma + \gamma(k_x^2 + k_y^2)a^2 \end{aligned}$$

which is free-electron-like, giving circular constant-energy surfaces near the centre of the Brillouin zone. If both  $k_x$  and  $k_y$  are close to  $\pi/a$ , write

$$k_x = \frac{\pi}{a} - \delta_x \quad k_y = \frac{\pi}{a} - \delta_y,$$

so that, remembering

$$\cos(a - b) = \cos(a) \cos(b) + \sin(a) \sin(b),$$

we have

$$\begin{aligned} E_{\mathbf{k}} &= -\alpha - 2\gamma(\cos(\pi - \delta_x a) + \cos(\pi - \delta_y a)) \\ &= -\alpha - 2\gamma(\cos(\pi) \cos(\delta_x a) - \sin(\pi) \sin(\delta_x a) \\ &\quad + \cos(\pi) \cos(\delta_y a) - \sin(\pi) \sin(\delta_y a)) \\ &= -\alpha + 2\gamma(\cos(\delta_x a) + \cos(\delta_y a)) \\ &= -\alpha + 4\gamma - \gamma(\delta_x^2 + \delta_y^2)a^2 \end{aligned}$$

giving circular constant-energy surfaces near the zone corners too.

Finally, in the middle of the band


$$\cos(k_x a) + \cos(k_y a) = 0,$$

the solutions to which are of the form

$$k_x a = \pi - k_y a,$$

or straight lines.

Finally, then, we have the constant energy surfaces for this tight-binding model.



### 7.6.3 Comments on tight binding theory

- Note that band width depends on two-centre integrals ( $\gamma$ ): for transition metals, this leads to narrow d-bands and wide s-bands.
- Near the top and bottom of bands, we have quadratic dependence on  $k$ .

A real band structure.

# Copper

